© 2026 International Conference on Multidisciplinary Sciences and Innovations . All Rights Reserved.

CLINICAL AND LABORATORY CHARACTERISTICS OF MULTIPLE ORGAN DYSFUNCTION SYNDROME IN INTENSIVE CARE

Adizova Madina Olimovna
Assistant Lecturer, Department of Clinical Sciences 1, Asia International University
Soliyev Alisher Uroqovich
Associate Professor, Department of Clinical Sciences 1, Asia International University

ABSTRACT

Multiple organ dysfunction syndrome represents a progressive and potentially reversible failure of two or more organ systems that cannot maintain homeostasis without therapeutic intervention. This syndrome develops as a consequence of severe infection, trauma, shock, or other critical conditions. The pathophysiology involves a dysregulated systemic inflammatory response characterized by excessive release of pro-inflammatory mediators, endothelial dysfunction, microvascular thrombosis, and impaired tissue perfusion. Clinical manifestations vary according to the affected organ systems, with respiratory, cardiovascular, renal, hepatic, hematological, neurological, and gastrointestinal dysfunction being most prominent. Laboratory findings include elevated creatinine and bilirubin levels, thrombocytopenia, elevated liver enzymes, metabolic acidosis, and coagulation abnormalities. The Sequential Organ Failure Assessment score remains the most widely utilized tool for quantifying organ dysfunction severity. Despite advances in critical care, mortality rates remain substantial, ranging from 30% to 100% depending on the number of failing organs. This thesis synthesizes current evidence regarding clinical and laboratory characteristics of multiple organ dysfunction syndrome in intensive care settings.

Key words: multiple organ dysfunction syndrome, intensive care, clinical manifestations, laboratory parameters, Sequential Organ Failure Assessment

INTRODUCTION

Multiple organ dysfunction syndrome is a clinical condition characterized by simultaneous or sequential impairment of function in two or more organ systems that are unable to maintain organismal homeostasis without therapeutic support . This syndrome represents a serious complication of critical illness and is frequently associated with sepsis, major trauma, hemorrhagic shock, and extensive burns . The syndrome was formally defined in 1991 during the American College of Chest Physicians and Society of Critical Care Medicine consensus conference, which introduced the concept of systemic inflammatory response syndrome as a unifying pathophysiological framework . The clinical significance of multiple organ dysfunction syndrome lies in its association with prolonged intensive care unit stays, increased healthcare resource utilization, and high mortality rates. Mortality increases substantially with the number of organs involved, ranging from approximately 10% for single organ failure to nearly 100% for failure of four or more organ systems . Among intensive care unit patients, sepsis remains the leading cause, affecting up to 30% of critically ill individuals . The economic burden extends beyond direct medical costs, encompassing long-term functional impairment in survivors and substantial strain on healthcare systems. This thesis aims to provide a comprehensive overview of the clinical and laboratory characteristics of multiple organ dysfunction syndrome in intensive care settings. The objective is to synthesize current evidence regarding organ-specific manifestations and diagnostic parameters that enable early recognition and severity assessment of this syndrome. Understanding these characteristics is essential for timely intervention and improved patient outcomes.

MAIN PART

© 2026 International Conference on Multidisciplinary Sciences and Innovations . All Rights Reserved.

Pathophysiological Basis and Etiological Factors - The pathogenesis of multiple organ dysfunction syndrome involves a complex interplay of dysregulated inflammatory responses, endothelial injury, microcirculatory disturbances, and cellular metabolic failure . An initially localized inflammatory response becomes systemic, leading to widespread endothelial activation and increased capillary permeability . Pro-inflammatory mediators including tumor necrosis factor alpha, interleukin-1, and interleukin-6 play central roles in propagating organ injury . Concurrently, reactive oxygen species contribute to oxidative stress and cellular damage, while disturbances in coagulation and fibrinolysis promote microvascular thrombosis . Infectious etiologies, particularly sepsis, represent the predominant cause of multiple organ dysfunction syndrome . Gram-negative bacteria, with Escherichia coli and Klebsiella species being most frequently isolated, account for a substantial proportion of cases . In tropical regions, parasitic infections including malaria constitute significant contributors . Non-infectious causes include severe trauma, extensive burns, acute pancreatitis, hemorrhagic shock, and ischemia-reperfusion injury following surgical procedures . The syndrome may also develop secondary to drug toxicity or thermal injury. Importantly, the type of inciting insult, patient comorbidities, and iatrogenic factors significantly influence both the development and progression of organ dysfunction .

Clinical Manifestations by Organ System

Respiratory Dysfunction. Respiratory failure is a hallmark feature of multiple organ dysfunction syndrome and frequently represents the initial organ system affected . Pulmonary dysfunction often progresses to acute respiratory distress syndrome, characterized by diffuse alveolar damage and increased pulmonary vascular permeability . This leads to impaired gas exchange manifesting as hypoxemia and hypercapnia. The ratio of arterial oxygen partial pressure to fractional inspired oxygen serves as the primary diagnostic parameter, with values below 150 indicating severe dysfunction . The majority of patients with multiple organ dysfunction syndrome require endotracheal intubation and mechanical ventilation . Clinical signs include progressive dyspnea, cyanosis, and refractory hypoxemia despite supplemental oxygen administration.

Cardiovascular Dysfunction. Cardiovascular manifestations encompass a spectrum of abnormalities including vasodilation, myocardial depression, and hypotension . Septic cardiomyopathy, characterized by impaired cardiac contractility and reduced stroke volume, is common among patients with sepsis-induced multiple organ dysfunction syndrome . Hypotension persisting despite adequate intravascular volume resuscitation indicates cardiovascular involvement . The clinical picture often reflects a mixed physiology of cardiogenic and distributive shock, complicating therapeutic management . Vasopressor and inotropic support requirements serve as quantitative indicators of cardiovascular dysfunction severity .

Renal Dysfunction. Acute kidney injury represents a frequent complication, with etiologies including ischemia-reperfusion injury, sepsis-induced renal dysfunction, and nephrotoxic insults . Clinical manifestations include oliguria or anuria, electrolyte abnormalities, and progressive azotemia. Serum creatinine concentration serves as the primary laboratory marker, with values exceeding 300 micromoles per liter indicating severe dysfunction . The need for renal replacement therapy, whether continuous or intermittent, signifies advanced renal failure . Hyperkalemia, metabolic acidosis, and volume overload often necessitate prompt initiation of dialysis.

Hepatic Dysfunction. Hepatic involvement in multiple organ dysfunction syndrome manifests through impaired synthetic function, coagulopathy, and hepatic encephalopathy . Reduced hepatic blood flow and hepatocellular injury contribute to impaired synthesis of both procoagulant and anticoagulant factors . Laboratory abnormalities include elevated serum bilirubin, transaminases, and alkaline phosphatase . Bilirubin levels exceeding 80 micromoles per liter with elevated serum

© 2026 International Conference on Multidisciplinary Sciences and Innovations . All Rights Reserved.

ammonia indicate severe dysfunction . Clinical signs may be insidious, often presenting with subtle findings such as cholestasis and mild transaminase elevation rather than overt jaundice .

Neurological Dysfunction. Neurological manifestations range from delirium to coma, with sepsis-associated encephalopathy being the most common presentation . The Glasgow Coma Scale serves as the primary assessment tool, with scores below 9 indicating severe dysfunction . Alterations in mental status or the onset of delirium often represent the first signs of decreased end-organ perfusion and incipient multiple organ dysfunction syndrome .

Gastrointestinal Dysfunction. Gastrointestinal involvement is characterized by mucosal ischemia, barrier dysfunction, and gut dysmotility . Compromised gut barrier integrity leads to bacterial translocation and systemic infection amplification . Clinical manifestations include ileus, delayed gastric emptying, and feeding intolerance . Nasogastric drainage exceeding 1000 milliliters per day indicates severe dysfunction . Gastrointestinal bleeding, acalculous cholecystitis, and pancreatitis represent additional manifestations included in some assessment systems .

Hematological Dysfunction. Hematological abnormalities include thrombocytopenia, disseminated intravascular coagulation, and anemia . Disseminated intravascular coagulation represents a consumptive coagulopathy characterized by widespread activation of coagulation and fibrinolysis, contributing to both microvascular thrombosis and bleeding complications . Platelet counts below 20,000 per cubic millimeter indicate severe dysfunction, while counts between 20,000 and 60,000 per cubic millimeter with prolonged prothrombin time and partial thromboplastin time suggest moderate involvement .

Laboratory Diagnostic Parameters and Scoring Systems - The diagnosis and severity assessment of multiple organ dysfunction syndrome rely on objective laboratory parameters and validated scoring systems. The Sequential Organ Failure Assessment score, introduced in 1996, remains the most widely utilized instrument for quantifying organ dysfunction . This score evaluates six organ systems: respiratory, renal, hepatic, hematological, cardiovascular, and neurological . Each system receives a score from zero to four based on the degree of dysfunction, with higher scores indicating greater severity. The pressure-adjusted heart rate, calculated as heart rate multiplied by the ratio of central venous pressure to mean arterial pressure, provides an alternative cardiovascular assessment parameter . Laboratory evaluations for suspected multiple organ dysfunction syndrome typically include complete blood cell counts with differential, comprehensive metabolic panels, coagulation studies, inflammatory markers, and arterial blood gas analysis . Elevated creatine phosphokinase exceeding 1000 units per liter, metabolic acidosis, and elevated liver enzymes are significantly associated with the presence of organ dysfunction . The Acute Physiology and Chronic Health Evaluation scoring system provides additional prognostic information . Emerging biomarkers including procalcitonin, presepsin, and various cytokines show promise for early diagnosis and prognostication, though their routine clinical application requires further validation

Management Implications - Management of multiple organ dysfunction syndrome requires a comprehensive approach directed at both the underlying cause and supportive care for failing organs . Prompt recognition and definitive treatment of the acute disease process remain fundamental priorities . Hemodynamic stabilization with appropriate fluid resuscitation and vasoactive support, adequate antimicrobial therapy for infectious etiologies, lung-protective mechanical ventilation, renal replacement therapy when indicated, and nutritional support constitute essential components of care . The concept of permissive physiological values compatible with survival, rather than strict normalization of parameters, has gained acceptance to minimize iatrogenic complications . Despite advances in critical care, multiple organ dysfunction syndrome continues to be associated with high mortality, underscoring the importance of early recognition and prevention strategies .

© 2026 International Conference on Multidisciplinary Sciences and Innovations . All Rights Reserved.

REFERENCES

1. Multiple organ dysfunction syndrome. Leksykon Wiedzy.
2. Rutledge and Sibbald Multiple Organ Dysfunction Assessment. In: Terapia Intensiva, Tomo 1. Generalidades.
3. Multiple Organ Dysfunction Syndrome: Molecular Mechanisms and Therapeutic Strategies. Signal Transduction and Targeted Therapy.
4. Niewydolność wielonarządowa. Leksykon Wiedzy.
5. Outcome definitions for Multiple Organ Dysfunction Syndrome. BMC Infectious Diseases supplementary material.
6. Multi-organ dysfunction syndrome. In: Elsevier Edited Volume.
7. Multiple Organ Failure: What You Need to Know. PMC.